



**QUEEN'S  
UNIVERSITY  
BELFAST**

## Label-Free Detection of Single-Base Mismatches in DNA by Surface-Enhanced Raman Spectroscopy

Papadopoulou, E., & Bell, S. E. J. (2011). Label-Free Detection of Single-Base Mismatches in DNA by Surface-Enhanced Raman Spectroscopy. *Angewandte Chemie International Edition*, 50(39), 9058-9061.  
<https://doi.org/10.1002/anie.201102776>

**Published in:**  
Angewandte Chemie International Edition

**Document Version:**  
Publisher's PDF, also known as Version of record

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)


**Publisher rights**  
Copyright © 2011 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

**General rights**  
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**  
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

# Label-Free Detection of Single-Base Mismatches in DNA by Surface-Enhanced Raman Spectroscopy\*\*

Evanthia Papadopoulou and Steven E. J. Bell\*



Angewandte  
Chemie

9058

WILEY  
ONLINE LIBRARY

© 2011 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Angew. Chem. Int. Ed. 2011, 50, 9058–9061

Detection of specific DNA sequences and identification of single-nucleotide polymorphisms (SNPs) are both important for diagnostic purposes, since alterations in the DNA sequence are the cause of the most common inherited disorders.<sup>[1]</sup> Methods for label-free identification of short DNA sequences are currently relatively limited. The most obvious choice is MALDI-TOF mass spectrometry, which is extensively used in SNP genotyping methods and has high sensitivity.<sup>[2]</sup> However, MALDI-TOF equipment is expensive, which partly offsets the significant advantage of being able to avoid labeling steps in the analysis. Surface-enhanced Raman spectroscopy (SERS) is an obvious alternative method, which has the potential to combine high sensitivity with low cost.<sup>[3]</sup> However, SERS studies of DNA most often involve detection of SERS labels;<sup>[4]</sup> even those that do not are typically carried out on sequences that have been thiolated to promote surface binding.<sup>[5]</sup> This approach not only increases the cost of the assay, but also the specific binding through a sulfur group gives the DNA a more tilted orientation that may prevent all the bases from interacting sufficiently with the substrate to provide good Raman signals. Indeed, it has been reported that the SERS spectra of thiolated DNA are completely dominated by adenine signals.<sup>[5]</sup> A recent approach utilized TERS (tip-enhanced Raman spectroscopy) to record the spectra of an unthiolated single poly C strand, in order to work towards a method of obtaining sequence information by moving the TERS probe in intervals of one base-to-base distance.<sup>[6]</sup> This approach, although promising, requires complex instrumentation, and, additionally, acquisition of Raman signals with single-base sensitivity is extremely challenging.


Herein we show that it is possible to obtain distinctive SERS spectra of unthiolated DNA sequences, which can be used to detect a single-nucleotide mismatch. This detection can be achieved by adsorbing the sequences nonspecifically through the nucleotide side chains onto the enhancing surface of silver colloids, where the sequences adopt a flattened configuration. To this end, the colloids were aggregated with an electrolyte that does not bind strongly in the silver surface ( $\text{MgSO}_4$  in this case),<sup>[7]</sup> and therefore allowed the DNA sequence to bind through its constituent bases. Previously, we demonstrated that silver colloids can provide very large SERS enhancements for the negatively charged mononucleotides when colloids were aggregated with  $\text{MgSO}_4$ .<sup>[7]</sup> We applied the same principle for the detection of single-base mismatches in short DNA sequences.

Initially, parallel studies were carried out using both thiolated and unthiolated 30-mer DNA-1 sequences (Table 1), which were either probed directly or after thermal pretreatment, as previously described.<sup>[5]</sup> All the samples gave

**Table 1:** DNA sequences used in this study.

Code	Sequence (5'–3')
DNA-1	GACTGCGACCAACCTAGCCTGCTATGATGT
HS-DNA-1	HS-CH <sub>2</sub> -DNA-1
DNA-2	CTT-TTT-CCT-GCA-TCC-TGT-CTG-GAA-A
DNA-3	CTT-TTT-CCT-GCA-TCC-TGT-CTG-GAA-G
DNA-4	ATA-AAT-CGC-CAT-TCG-TTG-ACT-AA
DNA-5	ATA-AAT-CGC-CAT-TCG-TTG-ACT-AC

acceptable SERS signals, as shown in Figure 1, thus demonstrating that thiolation is not necessary to obtain high-quality SERS spectra of DNA. In fact the unthiolated samples gave signals that were at least as large as the thiolated analogues. The spectra of the thermally treated thiolated sample strongly



**Figure 1.** SERS spectra of  $10^{-5}$  M solutions of a) thermally treated thiolated HS-DNA-1, b) untreated, thiolated HS-DNA-1, and c) unthiolated DNA-1. All spectra were recorded on citrate-reduced Ag colloids and aggregated with 0.1 M  $\text{MgSO}_4$ .

resemble the spectra of adenine,<sup>[8]</sup> which agrees with previous data obtained on Au substrates and was attributed to adenine having a stronger scattering cross-section than the other bases.<sup>[5]</sup> Interestingly, the SERS spectra of the unthiolated and thiolated form with no thermal pretreatment are different from the thermally treated sample, but similar to each other. This result suggests that the untreated thiolated sample adopts a similar adsorption configuration to the unthiolated sample, which is believed to lie flat with respect to the surface because of strong nonspecific binding between the bases and the metal surface. Similar binding interactions presumably also occur with thiolated strands<sup>[9]</sup> if they are not heat-treated. However, the most important point is that the spectra of simple unthiolated sequences are not dominated by the adenine modes, but show bands that arise from all the bases.

To aid the analysis of the DNA spectra, the SERS spectra of the three nucleobases poly A, poly C, and poly T were recorded and used as reference spectra. The SERS spectrum of dGMP is shown instead of poly G because poly G forms secondary structures through Hoogsteen base pairing.<sup>[10]</sup> We have found that these structures are not sufficiently flexible to

[\*] E. Papadopoulou, Dr. S. E. J. Bell  
Innovative Molecular Materials Group  
School of Chemistry & Chemical Engineering  
Queen's University, Belfast BT9 5AG (UK)  
E-mail: s.bell@qub.ac.uk


[\*\*] E. P. thanks Queen's University, Belfast for the award of a McClay Trust Studentship.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201102776>.

allow the nucleobases to directly adsorb to the surfaces of the Ag nanoparticles, and they therefore give only weakly enhanced Raman signals. For this work, hydroxylamine-reduced Ag colloids were used as the enhancing media because they are easier to synthesize than citrate-reduced colloids, and the low concentration of  $\text{Cl}^-$  ions introduced during their synthesis does not affect DNA binding. Indeed the SERS spectra (Figure S1 in the Supporting Information) show that when DNA is added to colloid aggregated with  $\text{MgSO}_4$ , it displaces the surface chloride layer and the  $244\text{ cm}^{-1}$  Ag–Cl band disappears. However, if the colloid is aggregated by using a high concentration of NaCl, no DNA Raman signals are observed because the DNA cannot compete with the high concentration of NaCl (0.1 M) for surface sites.


Irrespective of these considerations, the main point is that by using suitable conditions, the SERS spectra of unthiolated homopolymers or mononucleotides can be obtained (Figure 2); these spectra give characteristic bands that can be used for the identification of each nucleotide in a DNA sequence. For example, in the spectrum of the unthiolated form of DNA-1 (Figure 1c), the bands at 737 and  $682\text{ cm}^{-1}$  can be assigned to the ring breathing modes of poly A and poly G, respectively. The band at  $792\text{ cm}^{-1}$  can be assigned to the ring breathing modes of poly C and poly T, which overlap. The band at  $1635\text{ cm}^{-1}$  is assigned to the carbonyl stretching mode of poly C, while the intense  $1576\text{ cm}^{-1}$  poly G band is observed as a shoulder of the adenine band at  $1559\text{ cm}^{-1}$ .

Given that the SERS spectra of the unthiolated DNA strands show features associated with all the constituent bases, in principle it should be possible to detect even relatively small changes in the DNA sequence. We have



**Figure 2.** SERS spectra of  $10^{-6}\text{ M}$  solutions of the 24-mers a) poly A, b)  $10^{-5}$  dGMP, c) poly C, and d) poly T on hydroxylamine-reduced Ag colloid aggregated with 0.1 M  $\text{MgSO}_4$ .

tested the ability to detect A→G and C→A polymorphisms in 25-mer and 23-mer DNA sequences. Figure 3 shows the SERS spectra of DNA-2 and DNA-3, where the adenosine nucleotide in DNA-2 is replaced by a guanosine nucleotide in DNA-3. Most of the features appear unchanged but the band at  $1559\text{ cm}^{-1}$  in the DNA-2 spectrum is broader and shifts to  $1569\text{ cm}^{-1}$  in the DNA-3 spectrum. This observation is consistent with the strongest dGMP band, which lies at  $1576\text{ cm}^{-1}$  and increases in intensity in DNA-3.




**Figure 3.** SERS spectra of a) DNA-2 and b) DNA-3. c) Simple difference spectrum of DNA-3 minus DNA-2 ((b)–(a)) shown with its intensity expanded three times. d) Model difference spectrum, dGMP minus poly A. All spectra are for hydroxylamine-reduced Ag colloids aggregated with 0.1 M  $\text{MgSO}_4$ .

The best way to highlight the changes in the spectra is to digitally subtract them, which should remove the contributions from unchanged nucleotides to give difference spectra that contain positive and negative features corresponding to the exchanged nucleotides. Figure 3c shows the experimental difference spectrum. Negative bands that can be assigned to poly A vibrations are observed at  $1334\text{ cm}^{-1}$  and  $737\text{ cm}^{-1}$ . Positive G bands are observed at 1574 and  $1509\text{ cm}^{-1}$ , the expected increased band at  $1332\text{ cm}^{-1}$  is cancelled by the stronger negative band at  $1334\text{ cm}^{-1}$ , which corresponds to adenine. More strikingly, when the experimental difference spectrum is compared to a model spectrum created by subtracting the spectra of poly A from dGMP (Figure 3d), the agreement between the experimental and predicted pattern of band intensity changes is excellent. This result provides clear evidence that the changes in the spectra are due to the A→G polymorphism.

Each polymorphism would be expected to give a different pattern of band changes. For example, Figure 4 shows SERS data for DNA-4 and DNA-5, which correspond to C→A polymorphism. Again, small changes can be observed at the expected parts of the raw spectra, for example, the ratio of the





**Figure 4.** SERS spectra of a) DNA-4 and b) DNA-5. c) Simple difference spectrum of DNA-5 minus DNA-4 ((b)–(a)) shown with its intensity expanded eight times. d) Model difference spectrum, poly A minus poly C. All spectra are for hydroxylamine-reduced Ag colloids aggregated with 0.1 M  $\text{MgSO}_4$ .

intensity of the C band at  $792\text{ cm}^{-1}$  to that of the A band at  $737\text{ cm}^{-1}$  is slightly lower in the spectrum of ssDNA-5. However, the best evidence again comes from the difference spectra where the experimental data can be compared to the model obtained by subtracting the spectrum of poly C from that of poly A. Again there is excellent agreement between the expected and observed changes.

This work demonstrates that the SERS spectra of simple unthiolated DNA can be obtained under appropriate experimental conditions. These spectra show bands that are

characteristic of the constituent bases and are sufficiently reproducible that differences arising from single-base mismatch can be identified in short DNA strands.

### Experimental Section

All DNA oligomers were purchased from Eurogentec Ltd. (Belgium) and underwent sePOP (selective precipitation optimized process) desalting purification by the vendor. The unthiolated DNA sequences were dissolved in doubly distilled deionized water. The thiolated DNA-1 was thermally pretreated by heating the solutions to  $90^\circ\text{C}$  for 10–15 min and then rapid cooling in an ice bath. The SERS spectra were recorded on an Avalon Instrument RamanStation R1 (785 nm diode laser excitation). Typical exposure times were  $2 \times 60\text{ s}$ .

Received: April 21, 2011

Published online: July 19, 2011

**Keywords:** DNA · nanomaterials · nucleotides · RNA · SERS

- [1] A. C. Syvanen, *Nat. Rev. Genet.* **2001**, 2, 930–942.
- [2] J. Tost, I. G. Gut, *Clin. Biochem.* **2005**, 38, 335–350.
- [3] S. E. J. Bell, N. M. S. Sirimuthu, *Chem. Soc. Rev.* **2008**, 37, 1012–1024.
- [4] a) K. Faulds, F. McKenzie, W. E. Smith, D. Graham, *Angew. Chem.* **2007**, 119, 1861–1863; *Angew. Chem. Int. Ed.* **2007**, 46, 1829–1831; b) D. Graham, K. Faulds, *Chem. Soc. Rev.* **2008**, 37, 1042–1051.
- [5] A. Barhoumi, D. Zhang, F. Tam, N. J. Halas, *J. Am. Chem. Soc.* **2008**, 130, 5523–5529.
- [6] E. Bailo, V. Deckert, *Angew. Chem.* **2008**, 120, 1682–1685; *Angew. Chem. Int. Ed.* **2008**, 47, 1658–1661.
- [7] S. E. J. Bell, N. M. S. Sirimuthu, *J. Am. Chem. Soc.* **2006**, 128, 15580–15581.
- [8] E. Papadopoulou, S. E. J. Bell, *J. Phys. Chem. C* **2010**, 114, 22644–22651.
- [9] J. J. Storhoff, R. Elghanian, C. A. Mirkin, R. L. Letsinger, *Langmuir* **2002**, 18, 6666–6670.
- [10] K. Poon, R. B. Macgregor, *Biopolymers* **1998**, 45, 427–434.